

This listing of claims replaces all prior versions, and listings, of claims in the captioned application.

Listing of Claims:

1-10. (Previously Cancelled)

11. (Previously Withdrawn) A process of marking a receptor comprising the steps of a) radiolabelling a compound as defined in claim 1; b) administering said radiolabelled compound to biological material, c) detecting the emissions from the radiolabelled compound.


12. (Previously Withdrawn) A process of imaging an organ, characterized by, administering a sufficient amount of a radiolabelled compound of formula (I) in an appropriate composition, and detecting the emissions from the radioactive compound.

13. (Previously Cancelled)

14. (Previously Cancelled)

15-22. (Previously Cancelled)

23. A compound of formula



a *N*-oxide, a pharmaceutically acceptable addition salt or a stereochemically isomeric form thereof, wherein :

p represents an integer being 0, 1, or 2;

q represents an integer being 0, 1, or 2;

X represents O, S, NR³ or a direct bond;

R¹ represents hydrogen, hydroxy, halo, amino, C₁₋₆alkyl, C₁₋₆alkyloxy or mono- or di(C₁₋₄alkyl)aminoC₁₋₄alkylamino; in particular, hydrogen, methyl and hydroxy;

R² represents oxadiazolyl, thiazolyl, pyrimidinyl or pyridinyl; wherein said heterocycles each independently may optionally be substituted with one, or where possible, two or three substituents each independently selected from Het², R¹¹ and C₁₋₄alkyl optionally substituted with Het² or R¹¹;

each R⁴ independently represents C₁₋₆alkyl, halo, polyhaloC₁₋₆alkyl or C₁₋₆alkyloxy;

each R⁵ independently represents C₁₋₆alkyl, halo or C₁₋₆alkyloxy;

each R⁶ independently represents C₁₋₆alkylsulfonyl, aminosulfonyl or phenylC₁₋₄alkylsulfonyl;

each R⁷ and each R⁸ are independently selected from hydrogen, C₁₋₄alkyl, hydroxyC₁₋₄alkyl, dihydroxyC₁₋₄alkyl, aryl, arylC₁₋₄alkyl, C₁₋₄alkyloxyC₁₋₄alkyl, mono- or di(C₁₋₄alkyl)aminoC₁₋₄alkyl, arylaminocarbonyl, arylaminothiocarbonyl, C₃₋₇cycloalkyl, pyridinylC₁₋₄alkyl, Het³ and R⁶;

R⁹ and R¹⁰ are each independently selected from hydrogen, C₁₋₄alkyl, C₁₋₄alkylcarbonyloxyC₁₋₄alkylcarbonyl, hydroxyC₁₋₄alkylcarbonyl, C₁₋₄alkyloxyC₁₋₄alkylcarbonyl, Het³aminothiocarbonyl and R⁶;

each R¹¹ independently being selected from hydroxy, mercapto, cyano, nitro, halo, trihalomethyl, C₁₋₄alkyloxy, carboxyl, C₁₋₄alkyloxyC₁₋₄alkylcarbonyl, trihaloC₁₋₄alkylsulfonyloxy, R⁶, NR⁷R⁸, C(=O)NR⁷R⁸, aryl, aryloxy, arylcarbonyl, C₃₋₇cycloalkyl, C₃₋₇cycloalkyloxy, phthalimide-2-yl, Het³ and C(=O)Het³;

R¹² and R¹³ are each independently selected from hydrogen and C₁₋₄alkyl;

aryl represents phenyl optionally substituted with one, two or three substituents each independently selected from nitro, azido, halo, hydroxy, C₁₋₄alkyl, C₁₋₄alkyloxy, polyhaloC₁₋₄alkyl, NR⁹R¹⁰, R⁶, phenyl, Het³ and C₁₋₄alkyl substituted with NR⁹R¹⁰;

Het¹ represents a heterocycle selected from a heterocycle selected from imidazolyl, triazolyl, furanyl, oxazolyl, thiazolyl, thiazolinyl, thiadiazolyl, oxadiazolyl, pyridinyl, pyrimidinyl, pyrazinyl, piperidinyl, piperazinyl, triazinyl, benzothiazolyl, benzoxazolyl, purinyl, 1*H*-pyrazolo-[3,4-*d*]pyrimidinyl, benzimidazolyl, thiazolopyridinyl, oxazolopyridinyl, imidazo-[2,1-*b*]thiazolyl; wherein said heterocycles each independently may optionally be substituted with one, or where possible, two or three substituents each independently selected from Het², R¹¹ and C₁₋₄alkyl optionally substituted with Het² or R¹¹;

Het² represents furanyl, thienyl or pyridinyl; wherein said monocyclic heterocycles each independently may optionally be substituted with C₁₋₄alkyl;

Het³ represents pyrrolidinyl, piperidinyl, piperazinyl, morpholinyl, thiomorpholinyl; wherein said monocyclic heterocycles each independently may optionally be substituted with, where possible, one, two or three substituents each independently selected from C₁₋₄alkyl, C₁₋₄alkyloxy, C₁₋₄alkyloxycarbonyl, C₁₋₄alkylcarbonyl, phenylC₁₋₄alkyl, piperidinyl, NR¹²R¹³ and C₁₋₄alkyl substituted with NR¹²R¹³.

24. A compound according to claim 23 wherein the 6-azauracil moiety is in the para position relative to the central carbon atom.

25. A compound according to claim 24 wherein q is 1 or 2 and one R⁴ substituent is in the 4 position; and p is 1 or 2 and the one or two R⁵ substituents are in the ortho position relative to the central carbon atom.

26. A composition comprising a pharmaceutically acceptable carrier and, as active ingredient, a therapeutically effective amount of a compound as claimed in claim 23.

27. A process for preparing a composition as claimed in claim 26, wherein a pharmaceutically acceptable carrier is intimately mixed with a therapeutically effective amount of a compound as defined in claim 23.

28. **(Currently Cancelled)**

29. **(Currently Amended)** A method for treating one or more of: eosinophil-dependent inflammatory diseases bronchial asthma, atopic dermatitis, allergic-rhinitis or allergic conjunctivitis in a warm-blooded animal in need thereof comprising administering to the warm-blooded animal an effective amount of a compound of Claim 23.

30. **(Currently Cancelled).**

31. (New) A method for inhibiting IL-5 production in a warm-blooded animal, comprising administering to the warm-blooded animal an effective amount of a compound of claim 23.